

#### **Drug Facts**

#### Active ingredients

Purpose Naphazoline hydrochloride 0.025%....Redness reliever Pheniramine maleate 0.3%......Antihistamine

#### Uses

temporarily relieves itchy, red eyes due to:
• pollen • ragweed
• grass • animal hair and dander

#### Warnings

Do not use

• if you are sensitive to any ingredient in this

- Ask a doctor before use if you have

   heart disease high blood pressure

   narrow angle glaucoma
- trouble urinating due to an enlarged prostate gland

  When using this product

  • pupils may become enlarged temporarily

- do not touch tip of container to any surface to avoid contamination
- replace cap after each use
   remove contact lenses before using
   do not use if this solution changes color or
- becomes cloudy

  overuse may cause more eye redness

#### Stop use and ask a doctor if

- you feel eye pain
   changes in vision occur
   redness or irritation of the eye lasts
- condition worsens or lasts more than 72

EYE DROPS FORMERLY OCUHIST®

EYE ALLERGY RELIEF

EYE ALLERGY

RELIEF

ANTIHISTAMINE &

REDNESS RELIEVER

• ANTIHISTAMINE TO RELIEVE ITCHY EYES

Sterile.

## Drug Facts

(continued)

Keep out of reach of children If swallowed, get medical help or contact a Poison Control Center right away. Accidental swallowing by infants and children may lead to coma and marked reduction in

#### body temperature. Directions

- · adults and children 6 years of age and over: put 1 or 2 drops in the affected eye(s) up to four times a day
- for children under 6 years of age: consult a doctor

#### Other information

- some users may experience a brief tingling sensation
   store between 15° and 25°C (59° and 77°F)

# Inactive ingredients boric acid and sodium borate

buffer system preserved with benzalkonium chloride (0.01%) and edetate disodium (0.1%), sodium hydroxide and/or hydrochloric acid (to adjust pH), and purified water

### Questions? 1-800-723-7529, Monday to Friday, 9 AM - 5 PM EST

• RELIEVES REDNESS 1/2 FL. OZ. (15mL)

EYE ALLERGY RELIEF **Caution:** Do not use if Pfizer imprinted neckband on bottle is broken or missing. Pfizer

SM 38076018P

DISTRIBUTED BY: CONSUMER HEALTH CARE GROUP

PFIZER INC NEW YORK, NY 10017 www.prodhelp.com

EXP. DATE: LOT#

Jones CAD#4557A

8

# FORMERLY OCUHIST®

PLEASE READ CAREFULLY AND KEEP THIS INSERT FOR FUTURE REFERENCE

EYE ALLERGY

NTIHISTAMINE & REDNESS RELIEVER

EYE DROPS

#### **Drug Facts**

#### Active ingredients

**Purpose** 

Naphazoline hydrochloride 0.0259 Pheniramine maleate 0.3%......

.Antihistamine

#### Warnings

Do not use
• if you are sensitive to any ingredient in this product

#### Ask a doctor before use if you have

# **VISINE-A Insert**

Spec # SM11076022P

#### STANDARD FORMAT

Helvetica Condensed 100% Horizontal Scale

Drug Facts - 11 pt Helv Bold Italic

Drug Facts (continued) -

Drug Facts - 8 pt Helv Bold Italic

(continued) - 8 pt Helv Reg

Headings - 8 pt Helv Bold Italic

Sub Heads - 6 pt Helv Bold

Text - 6 pt Helv Reg, 1/2 pt leading

Round Bullets - 5 pt

1.5 pt Outline Box

1.5 pt Barline

0.5 pt Hairline

MM

3/64"- Gap (Border-Type)



# YOUR NEXT PURCHASE OF

NO EXPIRATION DATE

Antihistamine & Redness Reliever Eye Drops

CONSUMER: Limit one coupon per purchase specified. Any other use constitutes fraud. Void if copied, transferred or expired.

RETAILER: We will pay you for the face value of this coupon, plus & handling, if submitted in compliance with the Warner-Lambert Coupon Redemption Policy, available upon request. Void where prohibited, taxed, or restricted by law. Cash value 1/100¢. Mail coupons to: Warner-Lambert, P.O. Box #880142, El Paso TX 88588-0142. For questions regarding this coupon, call 1-800-648-0027. @2002 Pfizer Consumer Healthcare

\$1.00

#### Drug Facts (continued)

- When using this product

   pupils may become enlarged temporarily
- do not touch tip of container to any surface to avoid contamination
   replace cap after each use

- remove contact leases before using
   do not use if this solution changes color or becomes cloudy
   overuse may cause more eye redness

#### Stop use and ask a doctor if

- you feel eye pain
   changes in vision occur
   redness or irritation of the eye lasts
- condition worsens or lasts more than 72 hours

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Accidental swallowing by infants and children may lead to coma and marked reduction in body temperature.

#### Directions

- adults and children 6 years of age and over: put 1 or 2 drops in the affected eye(s) up to four times a day
   children under 6 years of age: consult a doctor

#### Other information

- use before the expiration date marked on the carton or bottle
   some users may experience a brief tingling sensation
   store between 15 and 25°C (59° and 77°F)

**Inactive ingredients** boric acid and sodium borate buffer system preserved with benzalkonium chloride (0.01%) and edetate disodium (0.1%), sodium hydroxide and/or hydrochloric acid (to adjust pH), and purified water. The solution has a pH of 5.5-6.5 and a tonicity of 245-305 mOsm/Kg.

Questions? call 1-800-223-0182, Monday to Friday, 9 AM - 5 PM EST

Dist: PFIZER CONSUMER HEALTHCARE Morris Plains, NJ 07950 USA ©2001 Pfizer www.prodhelp.com

SM 11076022P



This is a representation of an electronic record that was signed electronically a	nd
this page is the manifestation of the electronic signature.	

/s/

\_\_\_\_\_

Linda Katz

6/14/02 12:58:05 PM